

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
9 October 2003 (09.10.2003)

PCT

(10) International Publication Number
WO 03/082869 A1

(51) International Patent Classification⁷: **C07D 471/04,**
A61P 25/00, A61K 31/437

Darren, Peter [GB/GB]; Eisai London Research Laboratories Ltd, Bernard Katz Building, University College London, Gower Street, London WC1E 6BT (GB).

(21) International Application Number: **PCT/GB03/01115**

(74) Agents: **CORNISH, K., V., J. et al.;** Kilburn & Strode, 20 Red Lion Street, London WC1R 4PJ (GB).

(22) International Filing Date: **17 March 2003 (17.03.2003)**

(25) Filing Language: **English**

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(26) Publication Language: **English**

(30) Priority Data:
0207488.8 28 March 2002 (28.03.2002) GB
0300400.9 8 January 2003 (08.01.2003) GB

(71) Applicant (*for all designated States except US*): **EISAI LONDON RESEARCH LABORATORIES LIMITED** [GB/GB]; Bernard Katz Building, University College London, Gower Street, London WC1E 6BT (GB).

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **GRACZYK, Piotr** [PL/GB]; Eisai London Research Laboratories Ltd, Bernard Katz Building, University College London, Gower Street, London WC1E 6BT (GB). **NUMATA, Hirotoshi** [JP/JP]; Tsukuba Research Laboratories, Eisai Co. Ltd, 5-1-3 Tokodai, Tsukuba, Ibaraki 300-2635 (JP). **BHATIA, Gurpreet** [GB/GB]; Eisai London Research Laboratories Ltd, Bernard Katz Building, University College London, Gower Street, London WC1E 6BT (GB). **MEDLAND,**


Published:

— *with international search report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 03/082869 A1

(54) Title: **AZAINDOLES AS INHIBITORS OF C-JUN N-TERMINAL KINASES**



(I)

(57) Abstract: The present invention relates to novel 5-substituted 7-azaindole compounds of formula (I), their use in the inhibition of c-Jun N-terminal kinases, their use in medicine and particularly in the prevention and/or treatment of neurodegenerative disorders related to apoptosis and/or inflammation. The invention also provides processes for manufacture of said compounds, compositions containing them and processes for manufacturing such compositions.

AZAINDOLES AS INHIBITORS OF C-JUN N-TERMINAL KINASES

The present invention relates to novel 5-substituted 7-azaindole compounds, their use in the inhibition of c-Jun N-terminal kinases, their use in medicine and particularly in the prevention and/or treatment of neurodegenerative disorders related to apoptosis and/or inflammation. The invention also provides processes for manufacture of said compounds, compositions containing them and processes for manufacturing such compositions.

c-Jun N-terminal kinases (hereinafter referred to as "JNKs") are members of the mitogen-activated protein kinase (MAPK) family. JNKs are involved in response to various stimuli, including proinflammatory cytokines and environmental stress. JNKs, and JNK3 in particular, play an important role during apoptotic death of cells and therefore have been implicated in various disorders including stroke, traumatic brain injury and other neurodegenerative diseases such as Parkinson disease, Alzheimer disease and others. Since JNK activity is a physiological regulator of AP-1 transcriptional activity, JNK inhibitors are expected to reduce inflammatory response.

Apoptosis is a form of cell death in which the cell actively participates in its own destruction in a process involving a characteristic series of biochemical and morphological changes which are regulated by specific cell death genes. The apoptotic cell death is a process that has been observed in the developing mammalian nervous system. In mice, the inactivation by homologous recombination of genes that encode proteins that promote apoptosis, such as the caspase-3 or the Bax protein, prevents developmental neuronal cell death. The destruction of genes that encode cell death suppressors such as Bcl-x, leads to enhanced neuronal cell death. There is increasing evidence that apoptosis plays an important role in the pathology of acute and chronic neurodegenerative diseases. For example, in transgenic mice overexpressing the anti-apoptotic Bcl-2 protein in the nervous system there is a decrease in infarct volume following cerebral ischemia. Similarly, injection of the caspase inhibitor BAF reduces neuronal cell death following hypoxia/ischaemia in neonatal rats. Another

example is spinal muscular atrophy (a motor neurondisease) where loss of function mutations in the SMN gene is associated with the disease. Recent data has shown that the wild type SMN protein binds to Bcl-2 and co-operates with it to inhibit apoptosis. These results suggest that inhibitors of neuronal apoptosis
5 could be beneficial in the treatment of human neurodegenerative diseases. There is increasing evidence that neuronal apoptosis is an important pathological feature of stroke, traumatic brain injury and other neurodegenerative diseases. Therefore, pharmacotherapy using inhibitors of neuronal apoptosis may provide a therapeutic benefit in neurodegenerative conditions.

10

A number of groups have studied the mechanisms of neuronal cell death using *in vitro* cell culture systems and the results suggest that in some systems the transcription factor c-Jun is activated by the removal of survival signals and promotes cell death.

15

Antibodies specific for c-Jun protected NGF-deprived rat sympathetic neurones from apoptosis. Analogous neuroprotection due to expression of a c-Jun dominant negative mutant has been demonstrated, whereas overexpression of wild type c-Jun protein was sufficient to induce apoptosis in the presence of NGF. Estus and co-workers recently showed that an increase in c-Jun RNA
20 levels occurs in cortical neurones undergoing apoptosis after treatment with β -amyloid peptide (Estus *et al.*, 1997, J. Neurosci. 17, 7736-7745). It has also been shown that c-Jun is required for apoptosis in cerebellar granule neurones deprived of survival signals.

25

c-Jun is activated by JNKs, which phosphorylate its transcriptional activation domain. In humans there are three JNK genes : JNK1, JNK2 and JNK3. The RNAs encoding JNK1 and JNK2 are expressed in many tissues, including the brain, but JNK3 is restricted to the nervous system and to a smaller extent the
30 heart and testes.

JNKs are strongly activated in cellular responses to various stresses such as UV radiation, heat shock, osmotic shock, DNA-damaging agents, and

proinflammatory cytokines such as $\text{TNF}\alpha$, $\text{IL-1}\beta$ and others. Upstream regulators of the JNK pathway include kinases such as SEK1, MKK7 and MEKK1. There is evidence that Jun kinase activity is required for neuronal apoptosis *in vitro*. Overexpression of MEKK1 in sympathetic neurones
5 increased c-Jun protein levels and phosphorylation and induced apoptosis in the presence of NGF indicating that activation of the Jun kinase pathway can trigger neuronal cell death. The Jun kinase pathway has been shown to be necessary for the death of differentiated PC12 cells deprived of NGF. Furthermore, compound CEP-1347, which inhibits the c-Jun pathway (upstream of Jun kinase), protects
10 motor neurones against cell death induced by survival factor withdrawal.

In JNK3 homozygous (-/-) knockout mice, epileptic seizures and death of hippocampal CA3 neurones induced by injection of kainic acid is blocked. This indicates that JNK3 is involved in certain forms of neuronal cell death *in vivo*. It
15 is also a critical component of GluR6-mediated excitotoxicity. Furthermore, JNK3 (-/-) mice appear to develop normally and are viable suggesting that JNK3 is not essential for development or viability.


Strong nuclear JNK3 immunoreactivity in the brain CA1 neurones of patients
20 with acute hypoxia suggests that JNK3 is involved in hypoxia-related neurodegeneration. Transient hypoxia, may also trigger apoptosis through JNK signaling pathway in developing brain neurones.

Furthermore, JNK3 immunoreactivity is colocalized with Alzheimer disease-
25 affected neurones. Moreover JNK3 is related to neurofibrillary pathology of Alzheimer disease. In particular, JNK3 induces robust phosphorylation of amyloid precursor protein (APP) thus affecting its metabolism in disease state.

The present inventors have provided compounds which are inhibitors of c-Jun
30 N-terminal kinases.

The first aspect of the present invention relates to a compound of formula (I) as defined below:

4



5

wherein:

R stands for carbocyclyl, substituted carbocyclyl, heterocyclyl, or substituted heterocyclyl, wherein

10 the optionally substituted carbocyclyl or optionally substituted heterocyclyl group is optionally fused to an unsaturated, partially unsaturated or fully saturated five to seven membered ring containing zero to three heteroatoms,

each substitutable carbon atom in R, including the optional fused ring, is
 15 optionally and independently substituted by one or more of C_{1-12} alkyl, C_{2-12} alkenyl, carbocyclyl, or heterocyclyl, halogen, haloalkyl, OR^2 , SR^2 , NO_2 , CN , NR^2R^2 , NR^2COR^2 , $NR^2CONR^2R^2$, NR^2COR^2 , $NR^2CO_2R^2$, CO_2R^2 , COR^2 , $CONR^2R^2$, $S(O)_2R^2$, $SONH_2$, $S(O)R^2$, $SO_2NR^2R^2$, $NR^2S(O)_2R^2$, wherein each R^2 may be the same or different and is as
 20 defined below and wherein:

the C_{1-12} alkyl optionally incorporates one or two insertions selected from the group consisting of $-O-$, $-C(O)-$, $-N(R^2)-$, $-S(O)-$ and $-S(O)_2-$ wherein each R^2 may be the same or different and is as defined below;

25 the C_{1-12} alkyl, carbocyclyl, or heterocyclyl group is optionally substituted by one or more of halogen, haloalkyl, OR^2 , SR^2 , NO_2 , CN , NR^2R^2 , NR^2COR^2 , $NR^2CONR^2R^2$, NR^2COR^2 , $NR^2CO_2R^2$, CO_2R^2 , COR^2 , $CONR^2R^2$, $S(O)_2R^2$, $SONH_2$, $S(O)R^2$, $SO_2NR^2R^2$, $NR^2S(O)_2R^2$; wherein each R^2 may be the same or different and
 30 is as defined below and

the carbocyclyl, or heterocyclyl group is optionally substituted by one or more C_{1-12} alkyl,

each saturated carbon in the optional fused ring is further optionally and independently substituted by $=O$, $=S$, $=NNHR^2$, NNR^2R^2 , $=N-OR^2$,

5

$=\text{NNHCOR}^2$, $=\text{NNHCO}_2\text{R}^2$, $=\text{NNSO}_2\text{R}^2$, or $=\text{NR}^2$, wherein each R^2 may be the same or different and is as defined below; and

each substitutable nitrogen atom in R is optionally substituted by R^3 , COR^2 , SO_2R^2 or CO_2R^2 , wherein each R^2 and R^3 may be the same or different and is as defined below;

R^2 is hydrogen, C_{1-12} alkyl or aryl, optionally substituted by one or more of C_{1-4} alkyl, halogen, C_{1-4} haloalkyl, OR^4 , SR^4 , NO_2 , CN , NR^4R^4 , NR^4COR^4 , $\text{NR}^4\text{CONR}^4\text{R}^4$, NR^4COR^4 , $\text{NR}^4\text{CO}_2\text{R}^4$, CO_2R^4 , COR^4 , CONR^4_2 , $\text{S(O)}_2\text{R}^4$, SONH_2 , $\text{S(O)}\text{R}^4$, $\text{SO}_2\text{NR}^4\text{R}^4$, $\text{NR}^4\text{S(O)}_2\text{R}^4$, wherein the C_{1-12} alkyl group optionally incorporates one or two insertions selected from the group consisting of $-\text{O}-$, $-\text{N(R}^4)-$, $-\text{S(O)}-$ and $-\text{S(O)}_2-$, wherein each R^4 may be the same or different and is as defined below;

R^3 is C_{1-12} alkyl or aryl, optionally substituted by one or more of C_{1-4} alkyl, halogen, C_{1-4} haloalkyl, OR^4 , SR^4 , NO_2 , CN , NR^4R^4 , NR^4COR^4 , $\text{NR}^4\text{CONR}^4\text{R}^4$, NR^4COR^4 , $\text{NR}^4\text{CO}_2\text{R}^4$, CO_2R^4 , COR^4 , CONR^4_2 , $\text{S(O)}_2\text{R}^4$, SONH_2 , $\text{S(O)}\text{R}^4$, $\text{SO}_2\text{NR}^4\text{R}^4$, $\text{NR}^4\text{S(O)}_2\text{R}^4$, wherein the C_{1-12} alkyl group optionally incorporates one or two insertions selected from the group consisting of $-\text{O}-$, $-\text{N(R}^4)-$, $-\text{S(O)}-$ and $-\text{S(O)}_2-$, wherein each R^4 may be the same or different and is as defined below;

R^4 is hydrogen, C_{1-4} alkyl, or C_{1-4} haloalkyl;

with the proviso that when R is phenyl substituted with branched C_6 -alkyl ($-\text{CH}(\text{CH}_2-\text{CH}(\text{CH}_3)\text{CH}_3)-\text{CH}_2-$) incorporating two insertions $-(\text{CO})-$ and $-\text{NH}-$, the C_6 -alkyl group is not substituted with CN ;

and the pharmaceutically acceptable salts, and other pharmaceutically acceptable biohydrolyzable derivatives thereof, including esters, amides, carbamates, carbonates, ureides, solvates, hydrates, affinity reagents or prodrugs thereof.

For the avoidance of doubt, when a group as defined above contains two or more radicals, e.g. the radical R^2 , as for example in the groups $\text{SO}_2\text{NR}^2\text{R}^2$ and NR^2COR^2 , the radicals R^2 may be the same or different.

For the purposes of this invention, "alkyl" means a straight chain or branched alkyl radical of 1 to 12 carbon atoms, preferably 1 to 6 carbon atoms and most preferably 1 to 4 carbon atoms including but not limited to methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl etc. The term "alkenyl" means a straight chain or branched alkylenyl radical of 2 to 12 carbon atoms, preferably 2 to 6 carbon atoms and most preferably 2 to 4 carbon atoms, and containing one or more carbon-carbon double bonds and includes but is not limited to ethylene, n-propyl-1-ene, n-propyl-2-ene, isopropylene, etc. The term "alkynyl" means a straight chain or branched alkynyl radical of 2 to 12 carbon atoms, preferably 2 to 6 carbon atoms and most preferably 2 to 4 carbon atoms, and containing one or more carbon-carbon triple bonds and includes but is not limited to ethynyl, 2-methylethynyl etc. The term "cycloalkyl" means an saturated or partly unsaturated 3-12 membered cyclic alkyl group and includes but not limited to cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl etc. Cycloalkyl groups may be optionally substituted or fused to one or more aryl, heterocyclyl or cycloalkyl group. "Heterocycloalkyl" means a 3-12 membered saturated or partly unsaturated cycloalkyl containing one or more hetero atom selected from N, S and O. "Haloalkyl" means an alkyl radical substituted with one or more halide atoms for example $\text{CH}_2\text{CH}_2\text{Br}$, CF_3 or CCl_3 .

"Carbocyclyl" relates to a saturated, partly unsaturated or unsaturated 3-10 membered hydrocarbon ring, including cycloalkyl and aryl.

"Aryl" means an aromatic 3-10 membered hydrocarbon containing one ring or being fused to one or more saturated or unsaturated rings including but not limited to phenyl, naphthyl, anthracenyl or phenanthracenyl.

"Heteroaryl" means an aromatic 3-10 membered aryl containing one or more heteroatoms selected from N, O or S and containing one ring or being fused to one or more saturated or unsaturated rings and.

"Heterocyclyl" means a 3-10 membered ring system containing one or more heteroatoms selected from N, O or S and includes heteroaryl. The heterocyclyl

system can contain one ring or may be fused to one or more saturated or unsaturated rings; the heterocyclyl can be fully saturated, partially saturated or unsaturated and includes but is not limited heteroaryl and heterocarbocyclyl, e.g. cyclohexyl, phenyl, acridine, benzimidazole, benzofuran, benzothiophene, benzoxazole, benzothiazole, carbazole, cinnoline, dioxin, dioxane, dioxolane, 5 dithiane, dithiazine, dithiazole, dithiolane, furan, imidazole, imidazoline, imidazolidine, indole, indoline, indolizine, indazole, isoindole, isoquinoline, isoxazole, isothiazole, morpholine, naphthyridine, oxazole, oxadiazole, oxathiazole, oxathiazolidine, oxazine, oxadiazine, phenazine, phenothiazine, 10 phenoxazine, phthalazine, piperazine, piperidine, pteridine, purine, pyran, pyrazine, pyrazole, pyrazoline, pyrazolidine, pyridazine, pyridine, pyrimidine, pyrrole, pyrrolidine, pyrroline, quinoline, quinoxaline, quinazoline, quinolizine, tetrahydrofuran, tetrazine, tetrazole, thiophene, thiadiazine, thiadiazole, thiatriazole, thiazine, thiazole, thiomorpholine, thianaphthalene, thiopyran, 15 triazine, triazole, and trithiane;


Halogen means F, Cl, Br or I, preferably F.

R is preferably substituted with one or more of alkyl (e.g. methyl, ethyl or propyl), haloalkyl (preferably CF_3), halogen (e.g. F, Cl or Br, preferably F), OR^8 , 20 SR^8 , SOR^8 , $(\text{NR}^8)_2$, wherein R^8 is independently selected from hydrogen, C_{1-4} alkyl or haloalkyl and is preferably phenyl or naphthyl. When R is phenyl it is preferably substituted in the 4-(para) position, e.g. by $\text{NR}^6 \text{R}^6$, where R^6 stands independently for H or C_{1-4} alkyl.


25

Representative compounds according to the first aspect of the invention are illustrated below.


5




10



15




20



25

30



The compounds of the invention may be provided as a salt, preferably as a pharmaceutically acceptable salt of compounds of formula (I). Examples of pharmaceutically acceptable salts of these compounds include those derived from organic acids such as acetic acid, malic acid, tartaric acid, citric acid, lactic acid, oxalic acid, succinic acid, fumaric acid, maleic acid, benzoic acid, salicylic acid, phenylacetic acid, mandelic acid, methanesulphonic acid, benzenesulphonic acid and p-toluenesulphonic acid, mineral acids such as hydrochloric and sulphuric acid and the like, giving methanesulphonate, benzenesulphonate, p-toluenesulphonate, hydrochloride and sulphate, and the like, respectively or those derived from bases such as organic and inorganic bases. Examples of suitable inorganic bases for the formation of salts of compounds for this invention include the hydroxides, carbonates, and bicarbonates of ammonia, lithium, sodium, calcium, potassium, aluminium, iron, magnesium, zinc and the like. Salts can also be formed with suitable organic bases. Such bases suitable for the formation of pharmaceutically acceptable base addition salts with compounds of the present invention include organic bases which are nontoxic and strong enough to form salts. Such organic bases are already well known in the art and may include amino acids such as arginine and lysine, mono-, di-, or trihydroxyalkylamines such as mono-, di-, and triethanolamine, choline, mono-, di-, and trialkylamines, such as methylamine, dimethylamine, and trimethylamine, guanidine; N-methylglucosamine; N-methylpiperazine; morpholine; ethylenediamine; N-benzylphenethylamine; tris(hydroxymethyl) aminomethane; and the like.


Salts may be prepared in a conventional manner using methods well known in the art. Acid addition salts of said basic compounds may be prepared by dissolving the free base compounds according to the first or second aspects of the invention in aqueous or aqueous alcohol solution or other suitable solvents containing the required acid. Where a compound of the invention contains an acidic function, a base salt of said compound may be prepared by reacting said compound with a suitable base. The acid or base salt may separate directly or can be obtained by concentrating the solution e.g. by evaporation. The compounds of this invention may also exist in solvated or hydrated forms.

The invention also extends to a prodrug of the aforementioned compounds. A prodrug is any compound that may be converted under physiological conditions or by solvolysis to any of the compounds of the invention or to a pharmaceutically acceptable salt of the compounds of the invention. A prodrug may be inactive when administered to a subject but is converted *in vivo* to an active compound of the invention.

The compounds of the invention may contain one or more asymmetric carbon atoms and may exist in racemic and optically active forms. The first aspect of the invention covers all of these compounds.


In accordance with the second aspect of the present invention, the compound of the general formula (I):

15




can be made by hydrogenating a compound of the general formula (II):

25




in which R is as defined above and hal stands for a halogen atom, principally F or Cl, e.g. using in the presence of a suitable metal catalyst, such as e.g. palladium on activated carbon, and suitable amine such as e.g. triethylamine. The reaction can be run using a solution of compound (II) in a single solvent (e.g. alcohol, such as methanol or ethanol) or a mixture of solvents including e.g. an alcohol, dichloromethane, chloroform, etc.

12



5

The compound of the general formula (II) can be made by halogenating a compound of the general formula (III) in the 2 position, e.g.:




10

15

where R is as defined above, e.g. using $P(O)Cl_3$ at elevated temperature (about $100^\circ C$).

The compound of the general formula (III) can be made from 7-azaindole according to processes known in the art, see for example Glennon, K. C. *et al.* (WO00/56710) and Viaud, M.-C. *et al.* (EP0737685) and Cheung, M. *et al.* (WO99/21859). An example of a suitable scheme for the production of the compound of the general formula (III) is:

20



25

The third aspect of the present invention provides a composition comprising a compound of the general formula (I) as defined above in combination with a pharmaceutically acceptable carrier, diluent or excipient.

13

The composition may also comprise one or more additional active agent, such as an anti-inflammatory agent (for example a p38 inhibitor, glutamate receptor antagonist, or a calcium channel antagonist), a chemotherapeutic agent and/or an antiproliferative agent.

5

Suitable carriers and/or diluents are well known in the art and include pharmaceutical grade starch, mannitol, lactose, magnesium stearate, sodium saccharin, talcum, cellulose, glucose, sucrose, (or other sugar), magnesium carbonate, gelatin, oil, alcohol, detergents, emulsifiers or water (preferably sterile). The composition may be a mixed preparation of a composition or may be a combined preparation for simultaneous, separate or sequential use (including administration).

10

The composition according to the invention for use in the aforementioned indications may be administered by any convenient method, for example by oral (including by inhalation), parenteral, mucosal (e.g. buccal, sublingual, nasal), rectal or transdermal administration and the compositions adapted accordingly.

15

For oral administration, the composition can be formulated as liquids or solids, for example solutions, syrups, suspensions or emulsions, tablets, capsules and lozenges.

20

A liquid formulation will generally consist of a suspension or solution of the compound or physiologically acceptable salt in a suitable aqueous or non-aqueous liquid carrier(s) for example water, ethanol, glycerine, polyethylene glycol or an oil. The formulation may also contain a suspending agent, preservative, flavouring or colouring agent.

25

A composition in the form of a tablet can be prepared using any suitable pharmaceutical carrier(s) routinely used for preparing solid formulations. Examples of such carriers include magnesium stearate, starch, lactose, sucrose and microcrystalline cellulose.

30

14

A composition in the form of a capsule can be prepared using routine encapsulation procedures. For example, powders, granules or pellets containing the active ingredient can be prepared using standard carriers and then filled into a hard gelatin capsule; alternatively, a dispersion or suspension can be prepared using any suitable pharmaceutical carrier(s), for example aqueous gums, celluloses, silicates or oils and the dispersion or suspension then filled into a soft gelatin capsule.

Compositions for oral administration may be designed to protect the active ingredient against degradation as it passes through the alimentary tract, for example by an outer coating of the formulation on a tablet or capsule.

Typical parenteral compositions consist of a solution or suspension of the compound or physiologically acceptable salt in a sterile aqueous or non-aqueous carrier or parenterally acceptable oil, for example polyethylene glycol, polyvinyl pyrrolidone, lecithin, arachis oil or sesame oil. Alternatively, the solution can be lyophilised and then reconstituted with a suitable solvent just prior to administration.

Compositions for nasal or oral administration may conveniently be formulated as aerosols, drops, gels and powders. Aerosol formulations typically comprise a solution or fine suspension of the active substance in a physiologically acceptable aqueous or non-aqueous solvent and are usually presented in single or multidose quantities in sterile form in a sealed container, which can take the form of a cartridge or refill for use with an atomising device. Alternatively the sealed container may be a unitary dispensing device such as a single dose nasal inhaler or an aerosol dispenser fitted with a metering valve which is intended for disposal once the contents of the container have been exhausted. Where the dosage form comprises an aerosol dispenser, it will contain a pharmaceutically acceptable propellant. The aerosol dosage forms can also take the form of a pump-atomiser.

15

Compositions suitable for buccal or sublingual administration include tablets, lozenges and pastilles, wherein the active ingredient is formulated with a carrier such as sugar and acacia, tragacanth, or gelatin and glycerin.

- 5 Compositions for rectal or vaginal administration are conveniently in the form of suppositories (containing a conventional suppository base such as cocoa butter), pessaries, vaginal tabs, foams or enemas.

Compositions suitable for transdermal administration include ointments, gels,
10 patches and injections including powder injections.


Conveniently the composition is in unit dose form such as a tablet, capsule or ampoule.

- 15 In addition, the present invention provides a process for the manufacture of a composition according to the invention, as described above. The manufacture can be carried out by standard techniques well known in the art and involves combining a compound according to the first aspect of the invention and the pharmaceutically acceptable carrier or diluent. The composition may be in any
20 form including a tablet, a liquid, a capsule, and a powder or in the form of a food product, e.g. a functional food. In the latter case the food product itself may act as the pharmaceutically acceptable carrier.

The present invention provides a compound of the first aspect or a composition
25 of the third aspect for use in therapy/medicine.

The fourth aspect of the present invention relates to a compound of the general formula I as defined below

30



(I)

or a composition containing the compound, for use in inhibiting JNK

35 wherein:

R stands for carbocyclyl, substituted carbocyclyl, heterocyclyl, or substituted heterocyclyl, wherein

the optionally substituted carbocyclyl or optionally substituted heterocyclyl group is optionally fused to an unsaturated, partially unsaturated or fully saturated five to seven membered ring containing zero to three heteroatoms,

each substitutable carbon atom in R, including the optional fused ring, is optionally and independently substituted by one or more of C₁₋₁₂ alkyl, C₂₋₁₂ alkenyl, carbocyclyl, or heterocyclyl, halogen, haloalkyl, OR², SR², NO₂, CN, NR²R², NR²COR², NR²CONR²R², NR²COR², NR²CO₂R², CO₂R², COR², CONR²R², S(O)₂R², SONH₂, S(O)R², SO₂NR²R², NR²S(O)₂R², wherein each R² may be the same or different and is as defined below and wherein:

the C₁₋₁₂ alkyl optionally incorporates one or two insertions selected from the group consisting of -O-, -C(O)-, -N(R²)-, -S(O)- and -S(O)₂- wherein each R² may be the same or different and is as defined below;

the C₁₋₁₂ alkyl, carbocyclyl, or heterocyclyl group is optionally substituted by one or more of halogen, haloalkyl, OR², SR², NO₂, CN, NR²R², NR²COR², NR²CONR²R², NR²COR², NR²CO₂R², CO₂R², COR², CONR²R², S(O)₂R², SONH₂, S(O)R², SO₂NR²R², NR²S(O)₂R²; wherein each R² may be the same or different and is as defined below and

the carbocyclyl, or heterocyclyl group is optionally substituted by one or more C₁₋₁₂ alkyl,

each saturated carbon in the optional fused ring is further optionally and independently substituted by =O, =S, =NNHR², NNR²R², =N-OR², =NNHCOR², =NNHCO₂R², =NNSO₂R², or =NR², wherein each R² may be the same or different and is as defined below; and

each substitutable nitrogen atom in R is optionally substituted by R³, COR², SO₂R² or CO₂R², wherein each R² and R³ may be the same or different and is as defined below;

17

5 R^2 is hydrogen, C_{1-12} alkyl or aryl, optionally substituted by one or more of C_{1-4} alkyl, halogen, C_{1-4} haloalkyl, OR^4 , SR^4 , NO_2 , CN , NR^4R^4 , NR^4COR^4 , $NR^4CONR^4R^4$, NR^4COR^4 , $NR^4CO_2R^4$, CO_2R^4 , COR^4 , $CONR^4_2$, $S(O)_2R^4$, $SONH_2$, $S(O)R^4$, $SO_2NR^4R^4$, $NR^4S(O)_2R^4$, wherein the C_{1-12} alkyl group optionally incorporates one or two insertions selected from the group consisting of $-O-$, $-N(R^4)-$, $-S(O)-$ and $-S(O_2)-$, wherein each R^4 may be the same or different and is as defined below;

10 R^3 is C_{1-12} alkyl or aryl, optionally substituted by one or more of C_{1-4} alkyl, halogen, C_{1-4} haloalkyl, OR^4 , SR^4 , NO_2 , CN , NR^4R^4 , NR^4COR^4 , $NR^4CONR^4R^4$, NR^4COR^4 , $NR^4CO_2R^4$, CO_2R^4 , COR^4 , $CONR^4_2$, $S(O)_2R^4$, $SONH_2$, $S(O)R^4$, $SO_2NR^4R^4$, $NR^4S(O)_2R^4$, wherein the C_{1-12} alkyl group optionally incorporates one or two insertions selected from the group consisting of $-O-$, $-N(R^4)-$, $-S(O)-$ and $-S(O_2)-$, wherein each R^4 may be the same or different and is as defined below;

15 R^4 is hydrogen, C_{1-4} alkyl, or C_{1-4} haloalkyl;

and the pharmaceutically acceptable salts, and other pharmaceutically acceptable biohydrolyzable derivatives thereof, including esters, amides, carbamates, carbonates, ureides, solvates, hydrates, affinity reagents or prodrugs thereof.

20

All preferred options for R , R^2 , R^3 and R^4 and any substitutions or insertions thereof are as set out in the first aspect of the present invention. Preferred features of the composition of the fourth aspect are set out in the third aspect.

25 The compounds of the fourth aspect of the present invention are inhibitors of JNK, such as JNK1, JNK2, or JNK3. In particular, the compounds of the present invention are inhibitors of JNK3. Preferably, the compounds of the present invention inhibit JNK3 specifically.

30 One advantage of the compounds of the present invention is that they show a good stability to liver microsomes, at least when tested *in vitro* and hence are not rapidly metabolically removed from the body.

The compounds are therefore useful for conditions for which inhibition of JNK activity is beneficial. Thus, preferably, this aspect provides a compound of the general formula (I), or a composition containing a compound of formula (I) as defined in the fourth aspect of the present invention, as described above, for the prevention or treatment of a JNK-mediated disorder. The compounds of the general formula I may thus be used for the inhibition of JNK, more preferably for the inhibition of JNK3.

A "JNK-mediated disorder" is any disease or deleterious condition in which JNK plays a role. Examples include neurodegenerative disorder (including dementia), inflammatory disease, a disorder linked to apoptosis, particularly neuronal apoptosis, autoimmune disease, destructive bone disorder, proliferative disorder, cancer, infectious disease, allergy, ischemia reperfusion injury, heart attack, angiogenic disorder, organ hypoxia, vascular hyperplasia, cardiac hypertrophy, thrombin induced platelet aggregation and any condition associated with prostaglandin endoperoxidase synthase-2. The compounds of the present invention may be used for any of these JNK-mediated disorders.

The compounds of the fourth aspect of the present invention are particularly useful for the prevention or treatment of a neurodegenerative disorder. In particular, the neurodegenerative disorder results from apoptosis and/or inflammation. Examples of neurodegenerative disorders are: dementia; Alzheimer's disease; Parkinson's disease; Amyotrophic Lateral Sclerosis; Huntington's disease; senile chorea; Sydenham's chorea; hypoglycemia; head and spinal cord trauma including traumatic head injury; acute and chronic pain; epilepsy and seizures; olivopontocerebellar dementia; neuronal cell death; hypoxia-related neurodegeneration; acute hypoxia; glutamate toxicity including glutamate neurotoxicity; cerebral ischemia; dementia linked to meningitis and/or neurosis; cerebrovascular dementia; or dementia in an HIV-infected patient.

The neurodegenerative disorder may be a peripheral neuropathy, including mononeuropathy, multiple mononeuropathy or polyneuropathy. Examples of peripheral neuropathy may be found in diabetes mellitus, Lyme disease or

uremia; peripheral neuropathy caused by a toxic agent; demyelinating disease such as acute or chronic inflammatory polyneuropathy, leukodystrophies, or Guillain-Barré syndrome; multiple mononeuropathy secondary to a collagen vascular disorder (e.g. polyarteritis nodosa, SLE, Sjögren's syndrome); multiple
5 mononeuropathy secondary to sarcoidosis; multiple mononeuropathy secondary to a metabolic disease (e.g. diabetes or amyloidosis); or multiple mononeuropathy secondary to an infectious disease (e.g. Lyme disease or HIV infection).

10 The compounds of the invention can also be used to prevent or treat disorders resulting from inflammation. These include, for example, inflammatory bowel disorder, bronchitis, asthma, acute pancreatitis, chronic pancreatitis, allergies of various types, and possibly Alzheimer's disease. Autoimmune diseases which may also be treated or prevented by the compounds of the present invention
15 include rheumatoid arthritis, systemic lupus erythematosus, glomerulonephritis, scleroderma, chronic thyroiditis, Graves's disease, autoimmune gastritis, diabetes, autoimmune haemolytic anaemia, autoimmune neutropenia, thrombocytopenia, atopic dermatitis, chronic active hepatitis, myasthenia gravis, multiple sclerosis, ulcerative colitis, Crohn's disease, psoriasis or graft vs host disease.

20 A compound of the present invention may be administered simultaneously, subsequently or sequentially with one or more other active agent, such as an anti-inflammatory agent e.g. p38 inhibitor, glutamate receptor antagonist, calcium channel antagonist, a chemotherapeutic agent or an antiproliferative
25 agent. For example, for acute treatment, a p38 inhibitor may be administered to a patient prior to administering a compound of the present invention.

The compounds of the invention will normally be administered in a daily dosage regimen (for an adult patient) of, for example, an oral dose of between 1 mg and
30 2000 mg, preferably between 30 mg and 1000 mg, e.g. between 10 and 250 mg or an intravenous, subcutaneous, or intramuscular dose of between 0.1 mg and 100 mg, preferably between 0.1 mg and 50 mg, e.g. between 1 and 25 mg of the compound of the formula (I) or a physiologically acceptable salt thereof

calculated as the free base, the compound being administered 1 to 4 times per day. Suitably the compounds will be administered for a period of continuous therapy, for example for a week or more.

5 Accordingly the fifth aspect of the present invention relates to a method of treating or preventing a JNK-mediated disorder in an individual, which method comprises administering to said individual a compound as defined in the fourth aspect or a composition containing that compound. The active compound is preferably administered in a cumulative effective amount. The individual may
10 be in need of the treatment or prevention. Any of the JNK-mediated disorders listed above in relation to the fourth aspect may be the subject of treatment or prevention according to the fifth aspect. One or more other active agent may be administered to the individual simultaneously, subsequently or sequentially to administering the compound. The other active agent may be an anti-
15 inflammatory agent such as a p38 inhibitor, glutamate receptor antagonist, calcium channel antagonist, a chemotherapeutic agent or an antiproliferative agent, but is preferably p38 inhibitor for acute treatment.

The sixth aspect of the present invention provides the use of a compound of the
20 general formula (I) as defined in the fourth aspect of the invention in the manufacture of a medicament for the prevention or treatment of a JNK-mediated disorder. The medicament may be used for treatment or prevention of any of the JNK-mediated disorders listed above in relation to the fourth aspect. Again, the compound of the present invention may be administered simultaneously,
25 subsequently or sequentially with one or more other active agent, preferably a p38 inhibitor for acute treatment.

According to the seventh aspect of the present invention, there is also provided an assay for determining the activity of the compounds of the present invention,
30 comprising providing a system for assaying the activity and assaying the activity of the compound. Preferably the assay is for the JNK inhibiting activity of the compound, more preferably it is for the JNK3-specific inhibiting activity of the compounds. The compounds of the invention may be assayed *in vitro*, *in vivo*,

21


in silico, or in a primary cell culture or a cell line. *In vitro* assays include assays that determine inhibition of either the kinase activity or ATPase activity of activated JNK. Alternatively, *in vitro* assays may quantitate the ability of a compound to bind JNK and may be measured either by radiolabelling the compound prior to binding, then isolating the inhibitor/JNK complex and determining the amount of the radiolabel bound or by running a competition experiment where new inhibitors are incubated with JNK bound to known radioligands. An example of an assay which may be used is Scintillation Proximity Assay (SPA), preferably using radiolabelled ATP. Another example is ELISA. Any type or isoform of JNK may be used in these assays.

In a yet further aspect of the present invention, there is provided a method of inhibiting the activity or function of a JNK, particularly JNK3, which method comprises exposing a JNK to a compound or a composition of the first or fourth aspect of the present invention. The method may be performed in a research model, *in vitro*, *in silico*, or *in vivo* such as in an animal model. A suitable animal model may be a kainic acid model in rat or mice, traumatic brain injury model in rat, or MPTP in mice.


All features of each of the aspects apply to all other aspects *mutatis mutandis*.

Below, the present invention is illustrated using non-limiting examples.

EXAMPLES

Synthesis of example 5-substituted 7-azaindole derivative 7

Scheme 1


3,3-Dibromo-1,3-dihydro-pyrrolo[2,3-b]pyridin-2-one (2)

Technical (90%) pyridinium tribromide (220.4 g, 0.62 mol) was added portionwise over a period of 30 min to a stirred suspension of 7-azaindole (1, 27.13 g, 0.23 mol) in *t*-BuOH (1.36 L). The mixture was stirred at r.t for 3 h, and more pyridinium tribromide (73.3 g, 0.21 mol) was added in one portion. After additional stirring at r.t. for 2 h, the solvent was evaporated under reduced pressure. The residue was separated between water:AcOEt=1:1 (4.2 L). The aqueous layer was extracted with AcOEt (2x800 mL). Combined organic solutions were washed with water (2x500 mL), brine, dried (MgSO₄) and concentrated to dryness in vacuum. The residue was triturated with CH₂Cl₂ (1500 mL) for 20 min. The solid was filtered off, washed with CH₂Cl₂ (250 mL) and dried in vacuum to afford 2 (49.85 g, 75%) as yellow powder. ¹H NMR (400

23

MHz, DMSO- d_6) δ 7.16 (dd, $J = 7.4, 5.1$ Hz, 1H), 7.98 (dd, $J = 7.4, 1.5$ Hz, 1H), 8.19 (dd, $J = 5.1, 1.5$ Hz, 1H), 11.97 (bs, 1H).


3,3,5-Tribromo-1,3-dihydro-pyrrolo[2,3-b]pyridin-2-one (3)



Bromine (13.4 mL, 0.262 mol) was added dropwise over a period of 30 min to a cooled (ice bath) and stirred suspension of **2** (37.86 g, 0.131 mol) in water:*t*-BuOH=1:1 (1500 mL). Cooling bath was removed and the mixture was stirred at r.t. overnight. Then the solution was cooled to 15 °C and saturated aqueous solution of NaHCO₃ (278 mL) was added. A yellow suspension, which was formed, was concentrated in vacuum (bath temperature <32 °C) until about 1000 mL of condensate was collected. The solid was filtered off, washed with water (200 mL), and dried in vacuum to afford **3** (40.85 g, 85%) as tan powder.

10

5-Bromo-1,3-dihydro-pyrrolo[2,3-b]pyridin-2-one (4)



Zinc dust (34.0 g, 0.52 mol) was added in small portion to a stirred suspension of **3** (40.85 g, 0.111 mol) in glacial acetic acid (1000 mL) at such a rate that the temperature was maintained between 20-25 °C (strongly exothermic reaction; external ice bath cooling). Addition took about 20 min. Cooling bath was removed and stirring was continued at r.t. for 2 h. The solid was filtered off, washed with toluene (50 mL) and triturated with CH₂Cl₂:MeOH=4:1 (2.5 L). The solution was decanted off and treated with 1.0 M aqueous Na₂CO₃ solution (170 mL). After stirring for 1 h the two layers were separated. The organic layer was washed again with 1.0 M aqueous Na₂CO₃ solution (50 mL). The combined aqueous layers were extracted with CH₂Cl₂:MeOH=4:1 (10x100 mL). Combined organic solutions were dried with MgSO₄ (200 g) and concentrated. The residual

20


25

24

solid was dissolved in THF (2000 mL) and insoluble material was filtered off. The filtrate was concentrated to dryness in vacuum to afford **4** (16.93 g, 72%) as tan solid. ^1H NMR (400 MHz, DMSO- d_6) δ 3.57 (s, 2H), 7.75 (m, 1H), 8.14 (m, 1H), 11.13 (bs, 1H).

5


5-(3-Fluoro-phenyl)-1,3-dihydro-pyrrolo[2,3-b]pyridin-2-one (5)



A mixture of **4** (16.63 g, 78.5 mmol), 3-fluorophenylboronic acid (16.47 g, 117.7 mmol), Pd(PPh₃)₂Cl₂ (2.73 g, 6.60 mmol), LiCl (9.95 g, 0.23 mol), 1.0 M aqueous Na₂CO₃ solution (196 mL, 0.196 mol) in EtOH (470 mL) – toluene (470 mL) was refluxed overnight. More Pd(PPh₃)₂Cl₂ (1.30 g, 3.14 mmol) was added and reflux was continued for 24 h. The mixture was cooled, and the organic layer was separated and washed with brine (100 mL). The washings were combined with the aqueous layer and extracted with AcOEt (4x400 mL). Combined extracts were washed with brine, added to the organic layer and dried with MgSO₄. The solution was concentrated to dryness in vacuum to give 26.98 g of brown semisolid, which was triturated with ether:hexane=1:1 (2x500 mL). The residue was dried in vacuum to afford **5** (16.85 g, 94%) as tan solid, which was used in the next step without further purification. ^1H NMR (400 MHz, CDCl₃) δ 3.66 (s, 2H), 7.08 (dddd, J = 8.4, 8.2, 2.4, 0.9 Hz, 1H), 7.22 (ddd, J = 10.0, 2.4, 1.7 Hz, 1H), 7.30 (ddd, J = 8.1, 1.7, 0.9 Hz, 1H), 7.43 (ddd, J = 8.2, 8.1, 6.0 Hz, 1H), 7.69 (s, 1H), 8.36 (d, J = 2.1 Hz, 1H), 8.98 (bs, 1H).

2-Chloro-5-(3-fluoro-phenyl)-1H-pyrrolo[2,3-b]pyridine (6)


25



A suspension of **5** (16.52 g, 72.4 mmol) in neat P(O)Cl_3 (21.5 mL, 0.231 mol) was stirred at 100-105 °C for 4 h. The mixture was then cooled to r.t., diluted with *p*-xylene (100 mL) and concentrated to dryness in vacuum. The residue was separated between saturated aqueous NaHCO_3 – AcOEt. 10% aqueous solution of Na_2CO_3 was added to basify the aqueous layer to pH 9. Organic phase was separated and the aqueous layer was extracted with AcOEt (8x300 mL). Combined organic solutions were dried MgSO_4 , concentrated, and the residue was purified by silicagel chromatography (SGC) using CH_2Cl_2 :AcOEt as eluent in gradient to afford recovered starting material **5** (0.76 g, 5%). The desired product was then crystallized from acetone to afford **6** (10.06 g, 56%), thin tan needles. ^1H NMR (400 MHz, CDCl_3) δ 6.47 (s, 1H), 7.08 (tdd, $J = 8.1, 2.3, 1.5$ Hz, 1H), 7.33 (ddd, $J = 9.9, 2.3, 1.6$ Hz, 1H), 7.38-7.48 (m, 2H), 8.03 (d, $J = 2.1$ Hz, 1H), 8.53 (d, $J = 2.1$ Hz, 1H), 11.46 (bs, 1H).

15

5-(3-Fluoro-phenyl)-1H-pyrrolo[2,3-b]pyridine (**7**)



A mixture of chloride **6** (5.23 g, 21.3 mmol), 10% Pd/C (2.7 g), Et_3N (3.6 mL, 25.8 mmol) in THF:MeOH=5:1 (180 mL) was stirred under H_2 overnight. More 10% Pd/C (1.3 g) was added and stirring was continued for 3 h. Catalyst was removed by filtration and the solution was concentrated to dryness in vacuum. The residue was purified by SGC with CH_2Cl_2 :AcOEt as eluent in gradient (up to 20% AcOEt) to afford **7** (5.23 g, 88%), greenish powder. ^1H NMR (400 MHz, CDCl_3) δ 6.59 (dd, $J = 3.5, 2.0$ Hz, 1H), 7.04-7.10 (m, 1H), 7.33-7.37 (m, 1H),

7.40-7.48 (m, 3H), 8.14 (d, $J = 2.1$ Hz, 1H), 8.57 (d, $J = 2.1$ Hz, 1H), 9.91 (bs, 1H).

JNK1, JNK2, JNK3 - SPA assay

5

A typical assay for testing the activity of compounds to inhibit JNK1, JNK2 and JNK3 enzymes is as follows:

1. Compound is dissolved in DMSO to a convenient concentration and this is
10 diluted in 10% DMSO to a five times concentrate of the desired starting
concentration (frequently 1:100).
2. 10 μ l of 500 mM EDTA is added to alternative wells of the Opti-plate row,
which will receive kinase reaction plus DMSO. This creates the negative
control.
- 15 3. For the JNK2 and JNK3 assay, compounds are prepared in six 2-fold
dilutions with water and each concentration is tested in duplicate. For the
JNK1 assay compounds are prepared in four 5-fold dilutions with water
which are tested in triplicate. Controls are treated identically.
4. 20 μ l per well of each compound concentration is transferred to an Opti-
20 plate, in duplicate.
5. 30 μ l (JNK2/3 SPA) or 50 μ l (JNK1 SPA) of substrate solution (25 mM
HEPES pH 7.5, 10mM magnesium acetate with 3.33 μ M ATP (JNK2/3) or
2 μ M ATP (JNK1), approximately 7.5 kBq [γ -33P] ATP, GST-c-Jun, in
water) is added to each well.
- 25 6. 50 μ l (JNK2/3 SPA) or 30 μ l (JNK1 SPA) of kinase solution (JNK in 25 mM
HEPES pH 7.5, 10mM Mg Acetate) is added to each well.



Kinase	Kinase per well (μ g)	GST-c-Jun per well (μ g)
JNK1	0.25	1
JNK2	0.2	1.2
JNK3	0.16	1.2

7. The plate is incubated for 30 minutes at room temperature.
8. 100 μ l of bead/stop solution is added to each well (5 mg/ml glutathione-
PVT-SPA beads, 40 mM ATP in PBS).

9. Plates are sealed and incubated for 30 minutes at room temperature, centrifuged for 10 minutes at 2500g and counted.
10. The IC_{50} values are calculated as the concentration of the compound being tested at which the phosphorylation of c-Jun is decreased to 50% of the control value. Example IC_{50} values for the compounds of this invention are given in Table 1.

Examples of inhibitory potency against JNK3 kinase


10 Table 1. IC_{50} values for selected compounds against JNK3 kinase

<u>Compound Number</u>	<u>Structure</u>	JNK3 IC_{50} (μM)
1		1.2
2		<0.5

Claims

1. A compound of formula (I) as defined below:

5



wherein:

10

R stands for carbocyclyl, substituted carbocyclyl, heterocyclyl, or substituted heterocyclyl, wherein

15

the optionally substituted carbocyclyl or optionally substituted heterocyclyl group is optionally fused to an unsaturated, partially unsaturated or fully saturated five to seven membered ring containing zero to three heteroatoms,

20

each substitutable carbon atom in R, including the optional fused ring, is optionally and independently substituted by one or more of C₁₋₁₂ alkyl, C₂₋₁₂ alkenyl, carbocyclyl, or heterocyclyl, halogen, haloalkyl, OR², SR², NO₂, CN, NR²R², NR²COR², NR²CONR²R², NR²COR², NR²CO₂R², CO₂R², COR², CONR²R², S(O)₂R², SONH₂, S(O)R², SO₂NR²R², NR²S(O)₂R², wherein each R² may be the same or different and is as defined below and wherein:

25

the C₁₋₁₂ alkyl optionally incorporates one or two insertions selected from the group consisting of -O-, -C(O)-, -N(R²)-, -S(O)- and -S(O₂)- wherein each R² may be the same or different and is as defined below;

30

the C₁₋₁₂ alkyl, carbocyclyl, or heterocyclyl group is optionally substituted by one or more of halogen, haloalkyl, OR², SR², NO₂, CN, NR²R², NR²COR², NR²CONR²R², NR²COR², NR²CO₂R², CO₂R², COR², CONR²R², S(O)₂R², SONH₂, S(O)R², SO₂NR²R², NR²S(O)₂R²; wherein each R² may be the same or different and is as defined below and

the carbocyclyl, or heterocyclyl group is optionally substituted by one or more C₁₋₁₂ alkyl,

each saturated carbon in the optional fused ring is further optionally and independently substituted by =O, =S, =NNHR², NNR²R², =N-OR²,
 5 =NNHCOR², =NNHCO₂R², =NNSO₂R², or =NR², wherein each R² may be the same or different and is as defined below; and

each substitutable nitrogen atom in R is optionally substituted by R³, COR², SO₂R² or CO₂R², wherein each R² and R³ may be the same or different and is as defined below;

10 R² is hydrogen, C₁₋₁₂ alkyl or aryl, optionally substituted by one or more of C₁₋₄ alkyl, halogen, C₁₋₄ haloalkyl, OR⁴, SR⁴, NO₂, CN, NR⁴R⁴, NR⁴COR⁴, NR⁴CONR⁴R⁴, NR⁴COR⁴, NR⁴CO₂R⁴, CO₂R⁴, COR⁴, CONR⁴₂, S(O)₂R⁴, SONH₂, S(O)R⁴, SO₂ NR⁴R⁴, NR⁴S(O)₂R⁴, wherein the C₁₋₁₂ alkyl group optionally incorporates one or two insertions
 15 selected from the group consisting of -O-, -N(R⁴)-, -S(O)- and -S(O₂)-, wherein each R⁴ may be the same or different and is as defined below;

R³ is C₁₋₁₂ alkyl or aryl, optionally substituted by one or more of C₁₋₄ alkyl, halogen, C₁₋₄ haloalkyl, OR⁴, SR⁴, NO₂, CN, NR⁴R⁴, NR⁴COR⁴, NR⁴CONR⁴R⁴, NR⁴COR⁴, NR⁴CO₂R⁴, CO₂R⁴, COR⁴, CONR⁴₂, S(O)₂R⁴,
 20 SONH₂, S(O)R⁴, SO₂ NR⁴R⁴, NR⁴S(O)₂R⁴, wherein the C₁₋₁₂ alkyl group optionally incorporates one or two insertions selected from the group consisting of -O-, -N(R⁴)-, -S(O)- and -S(O₂)-, wherein each R⁴ may be the same or different and is as defined below;

R⁴ is hydrogen, C₁₋₄ alkyl, or C₁₋₄ haloalkyl; with the proviso that when R is phenyl substituted with branched C₆-alkyl (-CH(CH₂-CH(CH₃)(CH₃))-CH₂-) incorporating two insertions -(CO)-and-NH-, the C₆-alkyl group is
 25 not substituted with -CN;

and the pharmaceutically acceptable salts, and other pharmaceutically acceptable biohydrolyzable derivatives thereof, including esters, amides, carbamates,
 30 carbonates, ureides, solvates, hydrates, affinity reagents or prodrugs thereof.

2. A compound as claimed in claim 1, wherein R is an aryl or heteroaryl radical, optionally substituted with one or more of alkyl, haloalkyl, halogen,

30

OR^8 , S R^8 , SO R^8 , $(\text{NR}^8)_2$, wherein R^8 is independently selected from hydrogen, C_{1-4} alkyl or haloalkyl.

3. A compound as claimed in claim 1 wherein R is an optionally substituted
5 aryl, preferably phenyl or naphthyl.

4. A compound as claimed in claim 3, wherein R is phenyl substituted in the 4-(para) position.


10 5. A compound as claimed in claim 3 or claim 4, wherein R is phenyl substituted by $\text{NR}^6 \text{R}^6$, where R^6 stands independently for H or C_{1-4} alkyl.

6. A compound as claimed in claim 3 or claim 4, wherein R is substituted
15 aryl and the substituent is F, Cl or Br, preferably F; or haloalkyl, preferably CF_3 , or alkyl, preferably methyl, ethyl or propyl.


7. A compound as claimed in claim 1, which is one of the following:

31


5




10



15






8. A prodrug of a compound as defined in any of claims 1 to 7.

33

9. A process for the manufacture of any one or more of the compounds of any one of claims 1 to 7 which comprises hydrogenating a compound of the general formula (II):


5



10. in which R is as defined in claim 1 and hal stands for a halogen atom, e.g. using hydrogen and a catalyst such as Pd-C.

10. A process as claimed in claim 9, wherein the compound of the general formula (II) is made by halogenating a compound of the general formula (III) in the 2 position

15



20

where R is as defined above and hal stands for halogen, e.g. using $P(O)Cl_3$ at elevated temperature (about $100^\circ C$).

25 11. A composition comprising a compound as defined in any of claims 1-7 in combination with a pharmaceutically acceptable carrier, diluent or excipient.

12. A composition as claimed in claim 11 further comprising one or more other active agent.

30

13. A composition as claimed in claim 11 wherein the composition further comprises an anti-inflammatory agent, for example a p38 inhibitor.


14. A process for the manufacture of a composition as defined in any of claims 11-13, comprising combining a compound as defined in any of claims 1-7, and any additional active agent, with the pharmaceutically acceptable carrier or diluent.

5

15. A compound as defined in any of claims 1-7, or a composition as defined in any of claims 11-13, for use in therapy.

16. A compound of formula (I)

10



15

or a composition comprising a compound of formula (I), for inhibiting JNK, wherein:

20 R stands for carbocyclyl, substituted carbocyclyl, heterocyclyl, or substituted heterocyclyl, wherein

the optionally substituted carbocyclyl or optionally substituted heterocyclyl group is optionally fused to an unsaturated, partially unsaturated or fully saturated five to seven membered ring containing zero to three heteroatoms,

25 each substitutable carbon atom in R, including the optional fused ring, is optionally and independently substituted by one or more of C₁₋₁₂ alkyl, C₂₋₁₂ alkenyl, carbocyclyl, or heterocyclyl, halogen, haloalkyl, OR², SR², NO₂, CN, NR²R², NR²COR², NR²CONR²R², NR²COR², NR²CO₂R², CO₂R², COR², CONR²R², S(O)₂R², SONH₂, S(O)R², SO₂NR²R², NR²S(O)₂R², wherein each R² may be the same or different and is as defined below and wherein:

30

the C₁₋₁₂ alkyl optionally incorporates one or two insertions selected from the group consisting of -O-, -C(O)-, -N(R²)-, -S(O)-

and $-S(O_2)-$ wherein each R^2 may be the same or different and is as defined below;

the C_{1-12} alkyl, carbocyclyl, or heterocyclyl group is optionally substituted by one or more of halogen, haloalkyl, OR^2 , SR^2 , NO_2 , CN , NR^2R^2 , NR^2COR^2 , $NR^2CONR^2R^2$, NR^2COR^2 , $NR^2CO_2R^2$, CO_2R^2 , COR^2 , $CONR^2_2$, $S(O)_2R^2$, $SONH_2$, $S(O)R^2$, $SO_2NR^2R^2$, $NR^2S(O)_2R^2$; wherein each R^2 may be the same or different and is as defined below and

the carbocyclyl, or heterocyclyl group is optionally substituted by one or more C_{1-12} alkyl,

each saturated carbon in the optional fused ring is further optionally and independently substituted by $=O$, $=S$, $=NNHR^2$, NNR^2R^2 , $=N-OR^2$, $=NNHCOR^2$, $=NNHCO_2R^2$, $=NNSO_2R^2$, or $=NR^2$, wherein each R^2 may be the same or different and is as defined below; and

each substitutable nitrogen atom in R is optionally substituted by R^3 , COR^2 , SO_2R^2 or CO_2R^2 , wherein each R^2 and R^3 may be the same or different and is as defined below;

R^2 is hydrogen, C_{1-12} alkyl or aryl, optionally substituted by one or more of C_{1-4} alkyl, halogen, C_{1-4} haloalkyl, OR^4 , SR^4 , NO_2 , CN , NR^4R^4 , NR^4COR^4 , $NR^4CONR^4R^4$, NR^4COR^4 , $NR^4CO_2R^4$, CO_2R^4 , COR^4 , $CONR^4_2$, $S(O)_2R^4$, $SONH_2$, $S(O)R^4$, $SO_2NR^4R^4$, $NR^4S(O)_2R^4$, wherein the C_{1-12} alkyl group optionally incorporates one or two insertions selected from the group consisting of $-O-$, $-N(R^4)-$, $-S(O)-$ and $-S(O_2)-$, wherein each R^4 may be the same or different and is as defined below;

R^3 is C_{1-12} alkyl or aryl, optionally substituted by one or more of C_{1-4} alkyl, halogen, C_{1-4} haloalkyl, OR^4 , SR^4 , NO_2 , CN , NR^4R^4 , NR^4COR^4 , $NR^4CONR^4R^4$, NR^4COR^4 , $NR^4CO_2R^4$, CO_2R^4 , COR^4 , $CONR^4_2$, $S(O)_2R^4$, $SONH_2$, $S(O)R^4$, $SO_2NR^4R^4$, $NR^4S(O)_2R^4$, wherein the C_{1-12} alkyl group optionally incorporates one or two insertions selected from the group consisting of $-O-$, $-N(R^4)-$, $-S(O)-$ and $-S(O_2)-$, wherein each R^4 may be the same or different and is as defined below;

R^4 is hydrogen, C_{1-4} alkyl, or C_{1-4} haloalkyl;

36

and the pharmaceutically acceptable salts, and other pharmaceutically acceptable biohydrolyzable derivatives thereof, including esters, amides, carbamates, carbonates, ureides, solvates, hydrates, affinity reagents or prodrugs thereof.

5 17. A compound or a composition as defined in claim 16, for selectively inhibiting JNK3.

18. A compound or a composition as defined in claim 16, for use in the prevention or treatment of a JNK-mediated disorder.

10

19. A compound or a composition as claimed in claim 18, wherein the disorder is a neurodegenerative disorder (including dementia), inflammatory disease, a disorder linked to apoptosis, particularly neuronal apoptosis, autoimmune disease, destructive bone disorder, proliferative disorder, cancer,
15 infectious disease, allergy, ischemia reperfusion injury, heart attack, angiogenic disorder, organ hypoxia, vascular hyperplasia, cardiac hypertrophy, thrombin induced platelet aggregation and/or any condition associated with prostaglandin endoperoxidase synthase-2.

20 20. A compound or composition as claimed in claim 19, wherein the neurodegenerative disorder results from apoptosis and/or inflammation.

21. A compound or composition as claimed in claim 19 or claim 20, wherein the neurodegenerative disorder is: dementia; Alzheimer's disease; Parkinson's
25 disease; Amyotrophic Lateral Sclerosis; Huntington's disease; senile chorea; Sydenham's chorea; hypoglycemia; head and spinal cord trauma including traumatic head injury; acute and chronic pain; epilepsy and seizures; olivopontocerebellar dementia; neuronal cell death; hypoxia-related neurodegeneration; acute hypoxia; glutamate toxicity including glutamate
30 neurotoxicity; cerebral ischemia; dementia linked to meningitis and/or neurosis; cerebrovascular dementia; or dementia in an HIV-infected patient.

22. A compound or composition as claimed in claim 19 or 20, wherein the neurodegenerative disorder is a peripheral neuropathy, including mononeuropathy, multiple mononeuropathy or polyneuropathy, such as may be found in diabetes mellitus, Lyme disease or uremia; peripheral neuropathy
5 caused by a toxic agent; demyelinating disease such as acute or chronic inflammatory polyneuropathy, leukodystrophies or Guillain-Barré syndrome; multiple mononeuropathy secondary to a collagen vascular disorder (e.g. polyarteritis nodosa, SLE, Sjögren's syndrome); multiple mononeuropathy secondary to sarcoidosis; multiple mononeuropathy secondary to a metabolic
10 disease (e.g. diabetes or amyloidosis); or multiple mononeuropathy secondary to an infectious disease (e.g. Lyme disease or HIV infection).

23. A compound or composition as claimed in claim 18, wherein the disorder is inflammatory bowel disorder; bronchitis; asthma; acute pancreatitis; chronic
15 pancreatitis; allergies of various types; Alzheimer's disease; autoimmune disease such as rheumatoid arthritis, systemic lupus erythematosus, glomerulonephritis, scleroderma, chronic thyroiditis, Graves's disease, autoimmune gastritis, diabetes, autoimmune haemolytic anaemia, autoimmune neutropenia, thrombocytopenia, atopic dermatitis, chronic active hepatitis, myasthenia gravis, multiple sclerosis,
20 ulcerative colitis, Crohn's disease, psoriasis or graft vs host disease.

24. A method of treating or preventing a JNK-mediated disorder in an individual, which method comprises administering to said individual a compound or a composition as defined in claim 16.

25

25. A method as claimed in claim 24, wherein the individual is in need of the treatment or prevention of the disorder.

26. A method as claimed in claim 24 or 25, wherein the disorder is a
30 neurodegenerative disorder (including dementia), inflammatory disease, a disorder linked to apoptosis, particularly neuronal apoptosis, autoimmune disease, destructive bone disorder, proliferative disorder, cancer, infectious disease, allergy, ischemia reperfusion injury, heart attack, angiogenic disorder,

organ hypoxia, vascular hyperplasia, cardiac hypertrophy, thrombin induced platelet aggregation and/or any condition associated with prostaglandin endoperoxidase synthase-2.

5 27. A method as claimed in claim 26, wherein the neurodegenerative disorder results from apoptosis and/or inflammation.

28. A method as claimed in claim 26 or 27, wherein the neurodegenerative disorder is: dementia; Alzheimer's disease; Parkinson's disease; Amyotrophic
10 Lateral Sclerosis; Huntington's disease; senile chorea; Sydenham's chorea; hypoglycemia; head and spinal cord trauma including traumatic head injury; acute and chronic pain; epilepsy and seizures; olivopontocerebellar dementia; neuronal cell death; hypoxia-related neurodegeneration; acute hypoxia; glutamate toxicity including glutamate neurotoxicity; cerebral ischemia;
15 dementia linked to meningitis and/or neurosis; cerebrovascular dementia; or dementia in an HIV-infected patient.

29. A method as claimed in claim 26 or 27, wherein the neurodegenerative disorder is a peripheral neuropathy, including mononeuropathy, multiple
20 mononeuropathy or polyneuropathy, such as may be found in diabetes mellitus, Lyme disease or uremia; peripheral neuropathy caused by a toxic agent; demyelinating disease such as acute or chronic inflammatory polyneuropathy, leukodystrophies or Guillain-Barré syndrome; multiple mononeuropathy secondary to a collagen vascular disorder (e.g. polyarteritis nodosa, SLE,
25 Sjögren's syndrome); multiple mononeuropathy secondary to sarcoidosis; multiple mononeuropathy secondary to a metabolic disease (e.g. diabetes or amyloidosis); or multiple mononeuropathy secondary to an infectious disease (e.g. Lyme disease or HIV infection).

30 30. A method as claimed in claim 24, 25 or 26, wherein the disorder is inflammatory bowel disorder; bronchitis; asthma; acute pancreatitis; chronic pancreatitis; allergies of various types; Alzheimer's disease; autoimmune disease such as rheumatoid arthritis, systemic lupus erythematosus, glomerulonephritis,

scleroderma, chronic thyroiditis, Graves's disease, autoimmune gastritis, diabetes, autoimmune haemolytic anaemia, autoimmune neutropenia, thrombocytopenia, atopic dermatitis, chronic active hepatitis, myasthenia gravis, multiple sclerosis, ulcerative colitis, Crohn's disease, psoriasis or graft vs host disease.

5

31. A method as claimed in any of claims 24-30, wherein one or more other active agent is administered to the individual simultaneously, subsequently or sequentially to administering the compound.

10

32. A method as claimed in claim 31, wherein the other active agent is an anti-inflammatory agent such as a p38 inhibitor.

33. Use of a compound as defined in claim 16 in the manufacture of a medicament for the prevention or treatment of a JNK-mediated disorder.

15

34. Use as claimed in claim 33, wherein the disorder is a neurodegenerative disorder (including dementia), inflammatory disease, a disorder linked to apoptosis, particularly neuronal apoptosis, autoimmune disease, destructive bone disorder, proliferative disorder, cancer, infectious disease, allergy, ischemia reperfusion injury, heart attack, angiogenic disorder, organ hypoxia, vascular hyperplasia, cardiac hypertrophy, thrombin induced platelet aggregation and/or any condition associated with prostaglandin endoperoxidase synthase-2.

20

35. Use as claimed in claim 34, wherein the neurodegenerative disorder results from apoptosis and/or inflammation.

25

36. Use as claimed in claim 34 or 35, wherein the neurodegenerative disorder is: dementia; Alzheimer's disease; Parkinson's disease; Amyotrophic Lateral Sclerosis; Huntington's disease; senile chorea; Sydenham's chorea; hypoglycemia; head and spinal cord trauma including traumatic head injury; acute and chronic pain; epilepsy and seizures; olivopontocerebellar dementia; neuronal cell death; hypoxia-related neurodegeneration; acute hypoxia; glutamate toxicity including glutamate neurotoxicity; cerebral ischemia;

30

dementia linked to meningitis and/or neurosis; cerebrovascular dementia; or dementia in an HIV-infected patient.

37. Use as claimed in claim 34 or 35, wherein the neurodegenerative disorder is a peripheral neuropathy, including mononeuropathy, multiple mononeuropathy or polyneuropathy, such as may be found in diabetes mellitus, Lyme disease or uremia; peripheral neuropathy caused by a toxic agent; demyelinating disease such as acute or chronic inflammatory polyneuropathy, leukodystrophies or Guillain-Barré syndrome; multiple mononeuropathy secondary to a collagen vascular disorder (e.g. polyarteritis nodosa, SLE, Sjögren's syndrome); multiple mononeuropathy secondary to sarcoidosis; multiple mononeuropathy secondary to a metabolic disease (e.g. diabetes or amyloidosis); or multiple mononeuropathy secondary to an infectious disease (e.g. Lyme disease or HIV infection).

15

38. Use as claimed in claim 34, wherein the disorder is inflammatory bowel disorder; bronchitis; asthma; acute pancreatitis; chronic pancreatitis; allergies of various types; Alzheimer's disease; autoimmune disease such as rheumatoid arthritis, systemic lupus erythematosus, glomerulonephritis, scleroderma, chronic thyroiditis, Graves's disease, autoimmune gastritis, diabetes, autoimmune haemolytic anaemia, autoimmune neutropaenia, thrombocytopenia, atopic dermatitis, chronic active hepatitis, myasthenia gravis, multiple sclerosis, ulcerative colitis, Crohn's disease, psoriasis or graft vs host disease.

20

39. Use as claimed in any of claims 33-38, wherein the medicament further includes one or more other active agent.

25

40. Use as claimed in claim 39, wherein the other active agent is an anti-inflammatory agent such as a p38 inhibitor.

30

41. An assay for determining the activity of the compounds as defined in any of claims 1-7, comprising providing a system for assaying the activity and assaying the activity of a compound as defined in any of claims 1-7.

41

42. An assay as claimed in claim 41, wherein the assay is for the JNK inhibiting activity of the compound, preferably for the JNK3-specific inhibiting activity of the compound.

5

43. An assay as claimed in claim 41 or 42, wherein the assay is a Scintillation Proximity Assay (SPA) using radiolabelled ATP, or is ELISA.

10

44. A method of inhibiting the activity or function of a JNK, particularly JNK3, which method comprises exposing a JNK to a compound or composition as defined in claim 16.

45. A method as claimed in claim 44, which is performed in a research model.

15

46. A method as claimed in claim 45, wherein the research model is an animal model.

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/GB 03/01115

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D471/04 A61P25/00 A61K31/437

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 01 49288 A (MERCK FROSST CANADA INC ; ROBICHAUD JOEL STEPHANE (CA); ISABEL ELIS) 12 July 2001 (2001-07-12) page 115 -page 117; example A24	1-46
A	WO 01 47922 A (DEPRETS STEPHANIE ; LAI JUSTINE YEUN QUAI (GB); MORLEY ANDREW DAVID) 5 July 2001 (2001-07-05) the whole document	1-46
A	EP 1 106 621 A (FUJI PHOTO FILM CO LTD) 13 June 2001 (2001-06-13) the whole document	1-46



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

19 May 2003

Date of mailing of the international search report

27/05/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Baston, E

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/GB 03/01115

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0149288	A	12-07-2001	AU 2631401 A	16-07-2001
			CA 2396257 A1	12-07-2001
			EP 1248612 A1	16-10-2002
			WO 0149288 A1	12-07-2001
			US 2002052378 A1	02-05-2002
WO 0147922	A	05-07-2001	AU 2209401 A	09-07-2001
			BR 0017038 A	07-01-2003
			CA 2395593 A1	05-07-2001
			CN 1413213 T	23-04-2003
			CZ 20022207 A3	13-11-2002
			EP 1263759 A2	11-12-2002
			WO 0147922 A2	05-07-2001
			HU 0203895 A2	28-02-2003
			NO 20023032 A	21-06-2002
EP 1106621	A	13-06-2001	JP 2001163895 A	19-06-2001
			JP 2001163900 A	19-06-2001
			EP 1106621 A2	13-06-2001
			US 2003013088 A1	16-01-2003